Write your name here			
Surname	Other names		
Pearson Edexcel Level 1/Level 2 GCSE (9 - 1)	Candidate Number		
Combined Science Paper 3: Chemistry 1			
	Foundation Tier		
	Foundation Tier		
Paper 3: Chemistry 1	Foundation Tier		

Instructions

- Use **black** ink or ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must **show all your working out** with **your answer clearly identified** at the **end of your solution**.

Information

- The total mark for this paper is 60.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.
- In questions marked with an asterisk (*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.



Turn over 🕨



S50057A ©2016 Pearson Education Ltd.



(b	Some liquid is left in a warm room.	
	After a few days no liquid can be seen.	
	Give the name of the process that has occurred.	(1)
(c)	The freezing point of water is 0°C.	
	(i) Describe how the movement and arrangement of water particles changes when water is cooled from 10°C to −10°C.	(2)
	(ii) What is the structure of water?	(1)
	🖾 A ionic	
	B simple molecular (covalent)	
	C giant covalent	
	D metallic	
	(Total for Question 1 = 8 m	

2 Unreactive metals are found as uncombined metals in the Earth's crust. (a) Which of the following metals is found uncombined in the Earth's crust? (1) A aluminium B gold C sodium D zinc (b) When iron oxide is heated with carbon, iron is produced. (i) Complete the word equation for the reaction. (i) Complete the word equation for the reaction. (ii) What happens to the iron oxide during this reaction? A the iron oxide burns B the iron oxide is neutralised C the iron oxide is neutralised C the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO ₃ . In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed. (1)		
(1) A aluminium B gold C sodium D zinc (b) When iron oxide is heated with carbon, iron is produced. (i) Complete the word equation for the reaction. (ii) Complete the word equation for the reaction. (iii) What happens to the iron oxide during this reaction? A the iron oxide burns B the iron oxide is neutralised C the iron oxide is neutralised C the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO ₃ . In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide.	as uncombined metals in the Earth's crust.	
B gold C sodium D zinc (b) When iron oxide is heated with carbon, iron is produced. (i) Complete the word equation for the reaction. (i) Complete the word equation for the reaction. (2) iron oxide + carbon \rightarrow + (ii) What happens to the iron oxide during this reaction? (1) A the iron oxide burns B the iron oxide is neutralised C the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO ₃ . In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.	(1)	
□ C sodium □ D zinc (b) When iron oxide is heated with carbon, iron is produced. (i) Complete the word equation for the reaction. (2) iron oxide + carbon → (2) iron oxide + carbon → (2) (ii) What happens to the iron oxide during this reaction? (1) □ A the iron oxide burns □ B the iron oxide is neutralised □ C the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO ₃ . In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. $CuCO_3 → CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.		
□ D zinc (b) When iron oxide is heated with carbon, iron is produced. (i) Complete the word equation for the reaction. (i) Complete the word equation for the reaction. (2) iron oxide + carbon → + (ii) What happens to the iron oxide during this reaction? (1) □ A the iron oxide burns □ B the iron oxide is neutralised □ C the iron oxide is oxidised □ D the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO ₃ . In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.		
 (b) When iron oxide is heated with carbon, iron is produced. (i) Complete the word equation for the reaction. (2) iron oxide + carbon →		
(i) Complete the word equation for the reaction. (2) iron oxide + carbon \rightarrow +		
(2) iron oxide + carbon \rightarrow	d with carbon, iron is produced.	
iron oxide + carbon →	-	
(ii) What happens to the iron oxide during this reaction? (1) A the iron oxide burns B the iron oxide is neutralised C the iron oxide is oxidised D the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO ₃ . In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.		
(1) A the iron oxide burns B the iron oxide is neutralised C the iron oxide is oxidised D the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO ₃ . In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.		
Athe iron oxide burnsBthe iron oxide is neutralisedCthe iron oxide is oxidisedDthe iron oxide is reducedCcopper ore contains copper carbonate, $CuCO_3$.In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO , and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g.Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.	5	
 B the iron oxide is neutralised C the iron oxide is oxidised D the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO ₃ . In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. CuCO₃ → CuO + CO₂ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed. 		
 C the iron oxide is oxidised D the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO₃. In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. CuCO₃ → CuO + CO₂ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed. 		
 □ D the iron oxide is reduced (c) Copper ore contains copper carbonate, CuCO₃. In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. CuCO₃ → CuO + CO₂ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed. 		
(c) Copper ore contains copper carbonate, $CuCO_3$. In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.		
In the first stage of the extraction process, the copper carbonate is decomposed by heating to form copper oxide, CuO, and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.	educed	
by heating to form copper oxide, CuO, and carbon dioxide. $CuCO_3 \rightarrow CuO + CO_2$ When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.	per carbonate, CuCO ₃ .	
When 100 g of copper carbonate is decomposed completely in this way, it is found that the total mass of products is 100 g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.		
found that the total mass of products is 100g. Give a reason why the starting mass of copper carbonate is always the same as the mass of the products formed.	$CuCO_{3} \rightarrow CuO + CO_{2}$	
the mass of the products formed.		

DO NOT WRITE IN THIS AREA

(d) Zinc can be extracted from its ore by electrolysis or by heating the ore with carbon.

Give a reason for the method that is used.

(1)

(e) Figure 2 gives information about aluminium and tin.

metal	cost of 1 kg / £	amount in Earth's crust / %
aluminium	1.31	8
tin	12.60	0.0002

Figure 2

Give **two** reasons why it could be more important to recycle tin than to recycle aluminium. Use the information in Figure 2.

(2)

(Total for Question 2 = 8 marks)

Reason 1

OT WRITE IN THIS AREA

NOT WRITE IN THIS AREA

Reason 2

BLANK PAGE

Pearson Edexcel Level 1/Level 2 GCSE (9-1) in Combined Science – Sample Assessment Materials – Issue 1 – March 2016 © Pearson Education Limited 2016 3 An electrolysis experiment is carried out on different solutions, J, K and L.

Electricity is passed through each solution as shown in Figure 3.



Any products formed at the electrodes are identified.

The results are given in Figure 4.

solution	solution conducts electricity	product at cathode	product at anode
J	yes	copper	chlorine
К	no	none	none
L	yes	hydrogen	chlorine

Figure 4

(a) (i) State an improvement that can be made to the circuit to show that a current is flowing during the electrolysis.

(1)

DO NOT WRITE IN THIS AREA

	nat is meant by the term electro	lyte.			(2)
					(2)
(iii) Which o	f J , K and L are electrolytes?				
	J, R and L are electrolytes:				(1)
🖾 A Kon	ly				
🖾 B Jano	d L only				
🖾 C K an	d L only				
D J, K a	and L				
) Copper sulfa	ate solution was electrolysed for	r five minutes	using copper	electrodes.	
	•		song copper		
Figure 5 sho after electro	ows the mass of the anode and c				
	ows the mass of the anode and c				
	ows the mass of the anode and c	of the cathode	before electr		
	ows the mass of the anode and o olysis. mass of electrode	of the cathode anode	e before electro cathode		
	mass of electrode before electrodysis / g mass of electrode after electrolysis / g	of the cathode anode 1.16	cathode		
after electro	mass of electrode before electrodysis / g mass of electrode after electrolysis / g	of the cathode anode 1.16 0.85	cathode		
after electro	ows the mass of the anode and o olysis. mass of electrode before electrolysis / g mass of electrode after electrolysis / g Figu	of the cathode anode 1.16 0.85	cathode		(2)
after electro	ows the mass of the anode and o olysis. mass of electrode before electrolysis / g mass of electrode after electrolysis / g Figu	of the cathode anode 1.16 0.85	cathode		(2)
after electro	ows the mass of the anode and o olysis. mass of electrode before electrolysis / g mass of electrode after electrolysis / g Figu	of the cathode anode 1.16 0.85	cathode		(2)
after electro	ows the mass of the anode and o olysis. mass of electrode before electrolysis / g mass of electrode after electrolysis / g Figu	of the cathode anode 1.16 0.85	cathode		(2)
after electro	ows the mass of the anode and o olysis. mass of electrode before electrolysis / g mass of electrode after electrolysis / g Figu	of the cathode anode 1.16 0.85	cathode		(2)

(c)	Identify the products formed at the anode and cathode when molten
	potassium iodide is electrolysed.

(2)

Cathode

(d) In a different electrolysis, molten sodium fluoride is decomposed.

 $2NaF \rightarrow 2Na + F_2$

(relative masses: NaF = 42, Na = 23, $F_2 = 38$)

Calculate the maximum mass of sodium that could be formed from 168 g of sodium fluoride.

Anode

(2)

mass = g

(Total for Question 3 = 10 marks)

4 The apparatus in Figure 6 shows a piece of magnesium ribbon being heated.



Figure 6

During the heating, the magnesium reacts with oxygen from the air. The lid of the crucible was raised slightly from time to time. Magnesium oxide was formed as a white powder.

The experiment was repeated with different masses of magnesium.

The results are shown in Figure 7.

experiment	mass of magnesium used / g	mass of magnesium oxide formed / g	mass of oxygen in magnesium oxide / g
1	0.10	0.16	0.06
2	0.15	0.24	0.09
3	0.25	0.40	0.15
4	0.30	0.48	0.18
5	0.35	0.49	0.14
6	0.50	0.80	0.30

Figure 7

(a) (i) Draw a graph of the mass of oxygen in magnesium oxide against the mass of magnesium used. (3) mass of oxygen in magnesium oxide / g mass of magnesium used / g (ii) The result for experiment 5 is anomalous. The masses were all measured accurately. Suggest a reason for this anomalous result. (1) (b) Balance the equation for the reaction of magnesium with oxygen to form magnesium oxide. (1)Mg +O2 \rightarrow MgO

Calculate the relative formula mass of calcium nitrate, $Ca(NO_3)_2$. (relative atomic masses: Ca = 40, N = 14, O = 16)	(2)
relative formula ma	ss =
d) Two oxides of lead, R and S , were analysed.	
The empirical formula of oxide R was found to be PbO.	
The results of the analysis of oxide S showed it contained 0.207 g of lead combined with 0.032 g of oxygen.	
Show, by calculation, that the two oxides had different empirical formula (relative atomic masses: $O = 16$, $Pb = 207$)	e. (3)
(Total for Question 4	i = 10 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- 5 Substances can be pure or they can be mixtures.
 - (a) Which of these is a mixture?
 - 🖾 A chlorine
 - 🛛 B sodium
 - C sodium chloride
 - **D** sodium chloride solution
 - (b) Figure 8 shows some mixtures to be separated and possible methods of separation.

Place a tick (\checkmark) in one box in each row of the table to show the best method to separate the first named substance from each of the mixtures.

substance to	method of separation				
separate	crystallisation	filtration	simple distillation	fractional distillation	
sand from a mixture of sand and sodium chloride solution					
copper sulfate crystals from copper sulfate solution					
useful liquids from crude oil					

Figure 8

(3)



(ii) The chromatography paper, with the spot of mixture on it, was placed in a beaker with the bottom of the paper in water.

On Figure 10, complete the diagram showing the position of the chromatography paper with the spot of mixture at the start of the experiment.





(iii) The chromatography was carried out and the result is shown in Figure 11.





The blue spot had moved 14.5 cm and the solvent front had moved 15.3 cm

Calculate the R_f value of the substance in the blue spot, giving your answer to 2 significant figures.

 R_{f} value = $\frac{distance travelled by a dye}{distance travelled by solvent front}$

(2)

(1)

R_f value =

(d) **P**, **Q**, **R** and **S** are mixtures of food colourings.

They are investigated using paper chromatography.

Figure 12 shows the chromatogram at the end of the experiment.



Figure 12

(i) Which mixture contains an insoluble food colouring? (1)mixture **P** Α \mathbf{X} \mathbf{X} В mixture **Q** mixture **R** \mathbf{X} С D mixture S \times (ii) Give a change that could be made to the experiment to obtain an R_f value for the insoluble colouring. (1) (iii) Explain, by referring to Figure 12, which mixture is separated into the greatest number of soluble food colourings by this chromatography experiment. (2)

(Total for Question 5 = 12 marks)

- 6 Ionic compounds contain ions.
 - (a) The numbers of electrons, neutrons and protons in four particles, **W**, **X**, **Y** and **Z**, are shown in Figure 13.

particle	electrons	neutrons	protons
w	9	10	9
x	10	14	12
Y	16	16	16
Z	18	18	16

Figure 13

Explain which particle, **W**, **X**, **Y** or **Z**, is a negative ion.

(2)

(b) The electronic configurations of a lithium atom and of a fluorine atom are shown in Figure 14. Li F Figure 14 Lithium fluoride, LiF, is an ionic compound. It contains lithium cations and fluoride anions. Complete Figure 15 to show the electronic configurations and charges of the ions in lithium fluoride. (4) F charge on ion charge on ion Figure 15

*(c) Figure 16 shows the ability of different substances to conduct electricity.

substance	conducts electricity
solid calcium chloride	no
molten calcium chloride	yes
diamond	no
zinc	yes

Figure 16

Explain these results by referring to the structures of the substances.

(6)

(Total for Question 6 = 12 marks)

TOTAL FOR PAPER = 60 MARKS

81

DO NOT WRITE IN THIS AREA